

The Jacobian Conjecture as a Problem of Perturbative Quantum Field Theory

Abdelmalek Abdesselam

Département de Mathématiques

Université Paris XIII, Villetaneuse

Avenue J.B. Clément, F93430 Villetaneuse, France

February 1, 2008

Abstract

The Jacobian conjecture is an old unsolved problem in mathematics, which has been unsuccessfully attacked from many different angles. We add here another point of view pertaining to the so called formal inverse approach, that of perturbative quantum field theory.

Key words : Jacobian conjecture, Reversion, Quantum field theory.

I Introduction

The purpose of this modest note, for which we claim no originality except that of connecting apparently unrelated fields, is to draw the attention of theoretical physicists to one of the major unsolved problems of mathematics [35], viz. the Jacobian conjecture. The question is so simple that it was coined in [6] a problem in “high school algebra”. One can formulate it as follows.

Let $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a map written in coordinates as

$$F(x_1, \dots, x_n) = (F_1(x_1, \dots, x_n), \dots, F_n(x_1, \dots, x_n)) . \quad (1)$$

One says that F is a *polynomial map* if the functions $F_i : \mathbb{C}^n \rightarrow \mathbb{C}$ are polynomial. Suppose that the Jacobian determinant

$$JF(x_1, \dots, x_n) \stackrel{\text{def}}{=} \det \left(\frac{\partial F_i}{\partial x_j}(x_1, \dots, x_n) \right) \quad (2)$$

is *identically* equal to a nonzero constant. Show then that F is globally invertible (for the composition of maps) and that its inverse $G \stackrel{\text{def}}{=} F^{-1}$ is also a polynomial map.

Since it was first proposed in [25] (for $n = 2$ and polynomials with integral coefficients), this problem has resisted all attempts for a solution. In fact, this seemingly simple problem is quite an embarrassment. Indeed, some faulty proofs have even been published (see the indispensable [10] and [17] for a review). We will show here that the Jacobian conjecture can be formulated in very nice way as a question in preturbative quantum field theory (QFT). We also expect any future progress on this question to be beneficial not only for mathematics, but also for theoretical physics as it would enhance our understanding of perturbation theory.

Acknowledgments : The author is grateful to V. Rivasseau for early encouragements and collaboration on this project. Some of the ideas presented here are due to him. We thank D. Brydges, C. de Calan and J. Magnen for enlightening discussions. We also thank J. Feldman for his invitation to the Mathematics Department of the University of British Columbia where part of this work was done. The pictures in this article have been drawn using a software package kindly provided by J. Feldman.

II The formal inverse as a one-point correlation function

The most tempting, yet unfortunately least developed, line of attack on the Jacobian conjecture is the so called formal inverse approach. One tries to solve explicitly for $x = (x_1, \dots, x_n)$ in the equation $y = F(x)$, one then finds a power series expression for x in terms of $y = (y_1, \dots, y_n)$. By the uniqueness of the power series inverse, all one has to do then is to show that it is in fact a polynomial, that is the terms of high degree in the y variables vanish. One of the many reasons this approach is in its infancy is that it took

more than two centuries (say from [27] to [10]) to have a workable formula for the formal inverse in the multivariable case. Early contributions can be found in [28, 23, 15, 36, 29, 21]. An important contribution concerning formal inversion is due to Gurjar and Abhyankar [7]. Modern literature on reversion and Lagrange-Good type formulas is huge and we invite the reader to consult [10, 20, 22, 40] for more complete references. The first formula for the coefficients of the formal inverse power series G in terms of those of F is due to J. Towber and was first published in [41]. In physicists' terms ours is the following.

Claim : (A. A., V. Rivasseau) The formal solution of $y = F(x)$, without any assumption on F except that its linear part is invertible, is the perturbation expansion of the normalized one-point correlation function

$$x_i = \frac{1}{Z} \int_{\mathbb{C}^n} d\bar{\phi} d\phi \phi_i e^{-\bar{\phi}F(\phi) + \bar{\phi}y} \quad (3)$$

where $\bar{\phi}_1, \dots, \bar{\phi}_n, \phi_1, \dots, \phi_n$ are the components of a complex Bosonic field. The integration is over $\mathbb{C}^n$ with the measure

$$d\bar{\phi}d\phi \stackrel{\text{def}}{=} \prod_{i=1}^n \left(\frac{d(\text{Re } \phi_i) d(\text{Im } \phi_i)}{\pi} \right), \quad (4)$$

we used the notation $\bar{\phi}F(\phi) \stackrel{\text{def}}{=} \sum_{i=1}^n \bar{\phi}_i F_i(\phi_1, \dots, \phi_n)$, $\bar{\phi}y \stackrel{\text{def}}{=} \sum_{i=1}^n \bar{\phi}_i y_i$, and

$$Z \stackrel{\text{def}}{=} \int_{\mathbb{C}^n} d\bar{\phi}d\phi e^{-\bar{\phi}F(\phi) + \bar{\phi}y} \quad (5)$$

We obtained this expression by solving iteratively the equation $y = F(x)$ thereby generating a tree expansion in the same way one expresses the effective action $\Gamma(\phi)$ in terms of the logarithm $W(J)$ of the partition function in QFT (see [42] for instance). We then determined the Feynman rules of this tree expansion and finally the “path integral” formulation (3), only to realize that in fact our formula is closely related to the one introduced by G. Gallavotti, following a suggestion of G. Parisi, to express the Lindstedt perturbation series in the context of KAM theory [18].

A mathematician will undoubtedly shriek at the sight of equation (3). In the following, we will state and prove a precise theorem, using some analysis, for the case where $F_i(x) = x_i - H_i(x)$, with the $H_i(x)$, $1 \leq i \leq n$, being

homogenous of the same degree d . Indeed, it is enough to treat the cubic case $d = 3$, for all dimensions n , in order to prove the Jacobian conjecture in full generality [10]. However, formula (3) is completely combinatorial in nature and its proper setting is in the ring of formal power series with variables corresponding to the coefficients of F together with the y_i 's, over any field of zero characteristic. One simply has to define *formal Gaussian integration*, somewhat in the spirit of [8, 9]. We refer to the expository article [1] for a formulation and proof of our claim as a decent mathematical theorem. The latter article will also provide more details on how Feynman diagrams can be useful in algebraic combinatorics and how well they fit in the Joyal theory of combinatorial species [24]. We also refer to [2] for a very simple heuristic proof of the Lagrange-Good multivariable inversion formula, which becomes a fully rigorous and purely combinatorial proof when interpreted using the formalism of [1].

Now let $F_i(x) = x_i - H_i(x)$ with $H_i(x)$ written in *tensorial* notation as

$$H_i(x) = \frac{1}{d!} \sum_{j_1, \dots, j_d=1}^n w_{i,j_1 \dots j_d} x_{j_1} \dots x_{j_d} \quad (6)$$

so that the 1-contravariant and d -covariant tensor $w_{i,j_1 \dots j_d}$ is *completely symmetric* in the j indices. Let us write

$$\bar{\phi} w \phi^d \stackrel{\text{def}}{=} \sum_{i=1}^n \sum_{j_1, \dots, j_d=1}^n \bar{\phi}_i w_{i,j_1 \dots j_d} \phi_{j_1} \dots \phi_{j_d} \quad (7)$$

so that (3) becomes

$$G_i(y) = \frac{\int d\bar{\phi} d\phi \phi_i \exp \left(-\bar{\phi} \phi + \frac{1}{d!} \bar{\phi} w \phi^d + \bar{\phi} y \right)}{\int d\bar{\phi} d\phi \exp \left(-\bar{\phi} \phi + \frac{1}{d!} \bar{\phi} w \phi^d + \bar{\phi} y \right)} \quad (8)$$

The free propagator is represented as an oriented line

$$\begin{array}{c} \bullet \\ i \end{array} \longrightarrow \begin{array}{c} \bullet \\ j \end{array} = \delta_{ij} \quad (9)$$

for the contraction of a pair $\phi_i \bar{\phi}_j$. There are two types of vertices: the w -vertices represented by

$$\left. \begin{array}{c} \text{---} \bullet \text{---} \\ \text{---} \bullet \text{---} \\ \vdots \\ \text{---} \bullet \text{---} \end{array} \right\} d \text{ half lines} = \bar{\phi} w \phi^d \quad (10)$$

and the y -vertices represented by

$$\leftarrow \bullet = \overline{\phi} y . \quad (11)$$

As is well known in QFT, the numerator and denominator of (8) can be calculated by expanding

$$\exp \left(\frac{1}{d!} \overline{\phi} w \phi^d + \overline{\phi} y \right)$$

and integrating term by term with respect to the normalized complex Gaussian measure $d\overline{\phi} d\phi e^{-\overline{\phi}\phi}$. The result is a sum over all possible Feynman diagrams that can be built from the vertices of (9) and (10) (and the source ϕ_i for the numerator) by joining the half-lines of compatible directions. A quick look at the vertices shows that the only possible diagrams are trees connected to the source ϕ_i , or vacuum graphs made by an oriented loop of say $k \geq 1$ w -vertices to which $k(d-1)$ trees, whose leaves are y -vertices, are attached. When one factors out the denominator, the only diagrams that remain are made of a single tree with the source ϕ_i as its root. Therefore, at least formally, we have

$$G_i(y) = \sum_{V \geq 0} \sum_{N \geq 0} \frac{1}{V!N!} \sum_{\mathcal{T}} \mathcal{A}_i(\mathcal{T}) \quad (12)$$

where $\mathcal{T}$ is a Cayley tree (viewed as a set of unordered pairs) on a finite set $E = E(V, N)$. The latter is chosen, non canonically, once for each pair (V, N) , and must be the disjoint union of E_{root} of cardinality 1, E_{internal} of cardinality V and E_{leaf} of cardinality N . $\mathcal{T}$ is constrained by the condition that elements of $E_{\text{root}} \cup E_{\text{leaf}}$ have valence 1 while those of E_{internal} have valence $d+1$. This automatically enforces the relation $(d-1)V = N-1$ which can be checked by counting the half-lines. Even though we write, in the sequel, seemingly independent sums over V and N , the previous relation is always assumed.

We now define the *amplitude* $\mathcal{A}_i(\mathcal{T})$. One directs the edges of $\mathcal{T}$ towards the root in E_{root} . For each such edge $l \in \mathcal{T}$, one introduces an index i_l in the


set $\{1, \dots, n\}$. One then considers the expression $A_i(\mathcal{T}, (i_l)_{l \in \mathcal{T}})$ which is the product of the following factors.

- For each $a \in E_{\text{leaf}}$, if $l(a)$ is the unique line going from a , we take the factor $y_{i_{l(a)}}$.

- For each $b \in E_{\text{internal}}$, if $\{l_1(b), \dots, l_d(b)\}$ is the set of lines coming into b and $l_0(b)$ is the unique line leaving b , we take the factor $w_{i_{l_0(b)}, i_{l_1(b)} \dots i_{l_d(b)}}$.

The resulting monomial in the y 's and w 's is $A_i(\mathcal{T}, (i_l)_{l \in \mathcal{T}})$ by definition. Finally $\mathcal{A}_i(\mathcal{T})$ is the sum of $A_i(\mathcal{T}, (i_l)_{l \in \mathcal{T}})$ over all the indices $(i_l)_{l \in \mathcal{T}}$ except the index of the line arriving at the root which is fixed at the value i , the source index.

For example, with $d = 3$, the amplitude of the following tree with $V = 4$ and $N = 9$



is

$$\mathcal{A}_i(\mathcal{T}) = \sum_{\alpha_1, \dots, \alpha_{12}=1}^n w_{i, \alpha_1 \alpha_2 \alpha_3} w_{\alpha_1, \alpha_4 \alpha_5 \alpha_6} w_{\alpha_3, \alpha_7 \alpha_8 \alpha_9} w_{\alpha_9, \alpha_{10} \alpha_{11} \alpha_{12}} y_{\alpha_2} y_{\alpha_4} y_{\alpha_5} y_{\alpha_6} y_{\alpha_7} y_{\alpha_8} y_{\alpha_{10}} y_{\alpha_{11}} y_{\alpha_{12}} . \quad (13)$$

Note that, by the Cayley formula for the number of trees with preassigned valences, the sum over $\mathcal{T}$ has

$$\frac{((1 + V + N) - 2)!}{(1 - 1)!((d + 1) - 1)!^V (1 - 1)!^N} = \frac{(V + N - 1)!}{d!^V} \text{ terms.} \quad (14)$$

Let us introduce the norms

$$\|w\|_{\infty,1} \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} \sum_{j_1, \dots, j_d=1}^n |w_{i,j_1 \dots j_d}| \quad (15)$$

and

$$\|y\|_{\infty} \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} |y_i| . \quad (16)$$

We now have

Theorem 1 *The series*

$$G_i(y) = \sum_{V \geq 0} \sum_{N \geq 0} \frac{1}{V!N!} \sum_{\mathcal{T}} \mathcal{A}_i(\mathcal{T}) \quad (17)$$

is absolutely convergent, provided

$$\|y\|_{\infty} < R \stackrel{\text{def}}{=} \left(\frac{d!}{2^d \|w\|_{\infty,1}} \right)^{d-1} \quad (18)$$

and satisfies, on this domain of convergence,

$$\|G(y)\|_{\infty} \leq \frac{\|y\|_{\infty}}{1 - \frac{\|y\|_{\infty}}{R}} \quad (19)$$

and

$$F(G(y)) = y . \quad (20)$$

Proof : One easily proves by bounding the w and y factors in $A_i(\mathcal{T}, (i_l)_{l \in \mathcal{T}})$ by their moduli and summing the indices, starting with the leaves and progressing towards the root, that

$$|\mathcal{A}_i(\mathcal{T})| \leq \|w\|_{\infty,1}^V \|y\|_{\infty}^N \quad (21)$$

for any fixed tree $\mathcal{T}$. Therefore

$$\begin{aligned} & \sum_{V,N \geq 0} \frac{1}{V!N!} \sum_{\mathcal{T}} |\mathcal{A}_i(\mathcal{T})| \\ & \leq \sum_{V \geq 0} \frac{(dV)! \|w\|_{\infty,1}^V \|y\|_{\infty}^{(d-1)V+1}}{V!((d-1)V+1)!d!^V} \end{aligned} \quad (22)$$

$$\leq \|y\|_{\infty} \sum_{V \geq 0} \frac{(dV)!}{V!((d-1)V)!} \left(\frac{\|w\|_{\infty,1} \|y\|_{\infty}^{d-1}}{d!} \right)^V \quad (23)$$

and one simply uses

$$\frac{(dV)!}{V!((d-1)V)!} \leq 2^{dV} \quad (24)$$

to conclude the convergence proof and obtain the bound (19). Now observe that, on the convergence domain

$$G_i(y) = y_i + \sum_{\substack{V_i, N_i, \mathcal{T} \\ V \geq 1}} \frac{1}{V!N!} \mathcal{A}_i(\mathcal{T}) \quad (25)$$

where the last sum is over trees with at least one w -vertex linked directly to the root. This sum can be performed in the following way. One chooses, among the V internal w -vertices, the vertex $w_0 \in E_{\text{internal}}$ which hooks to the root. This costs a factor V . Then one divides the remaining vertices into an unordered collection of sets $E_1, \dots, E_d$ such that E_i has V_i w -vertices and N_i y -vertices. This costs a factor

$$\frac{1}{d!} \frac{(V-1)!}{V_1! \dots V_d!} \frac{N!}{N_1! \dots N_d!}.$$

Finally one sums over all possible trees $\mathcal{T}_1, \dots, \mathcal{T}_d$ on $E_1 \cup \{w_0\}, \dots, E_d \cup \{w_0\}$ as before. The corresponding amplitudes do not depend on the location of the sets E_i in E , but only on the cardinalities V_i and N_i . Therefore

$$G_i(y) = y_i + \sum_{\substack{V_1, \dots, V_d \geq 0 \\ N_1, \dots, N_d \geq 1}} \sum_{i_1, \dots, i_d=1}^n \sum_{\mathcal{T}_1, \dots, \mathcal{T}_d} \frac{1}{V!N!} \frac{V(V-1)!N!}{d!V_1! \dots V_d!N_1! \dots N_d!} w_{i, i_1 \dots i_d} \mathcal{A}_{i_1}(\mathcal{T}_1) \dots \mathcal{A}_{i_d}(\mathcal{T}_d) \quad (26)$$

$$= y_i + \sum_{i_1, \dots, i_d=1}^n \frac{1}{d!} w_{i, i_1 \dots i_d} G_{i_1}(y) \dots G_{i_d}(y) \quad (27)$$

$$= y_i + H_i(G(y)) \quad (28)$$

from which (20) follows. ■

As a result the Taylor series of G at the origin is the right compositional inverse of F . Now algebraic combinatorialists might not be too impressed by this since one can readily rewrite formula (12) under the form given by

Towber [40] or Singer [34]. So the series expansion of the formal inverse itself is not new. To obtain a real improvement on previous approaches one has to return to the more fundamental equation (3) and really consider the “integrals” appearing in it as, well, *integrals* on which one can try all the tools of ordinary calculus: integration by parts, change of variables... For an example of the mathematical utility of this way of proceeding, see [8, 9, 32]. **Remark :** Note that the generalized forest formula of Towber [40] can be easily derived from the perturbation expansion of higher correlation functions $\langle \phi_1^{\alpha_1} \dots \phi_n^{\alpha_n} \rangle$, where we used the standard statistical mechanics notation

$$\langle \Omega(\bar{\phi}, \phi) \rangle \stackrel{\text{def}}{=} \frac{1}{Z} \int d\bar{\phi} d\phi \Omega(\bar{\phi}, \phi) e^{-\bar{\phi}\phi + \frac{1}{d!} \bar{\phi} w \phi^d + \bar{\phi} y} . \quad (29)$$

III Comments on the Jacobian conjecture

III.1 What does the constant Jacobian condition mean?

Suppose that

$$F_i(x) = x_i - \sum_{j_1, \dots, j_d=1}^n \frac{1}{d!} w_{i,j_1 \dots j_d} x_{j_1} \dots x_{j_d} \quad (30)$$

is such that $JF(x) = 1$ for all x . Several conclusions can be drawn from this constraint. One that is due to V. Rivasseau is that our QFT model is *self-normalized*. In other words

$$Z = 1 . \quad (31)$$

Indeed, by writing the Feynman diagram expansion of

$$Z = \int d\bar{\phi} d\phi e^{-\bar{\phi}\phi + \frac{1}{d!} \bar{\phi} w \phi^d + \bar{\phi} y} \quad (32)$$

one can easily show that

$$\log Z = \sum_{k \geq 1} \frac{1}{k} \text{tr} [M(G(y))^k] \quad (33)$$

where $M(x)$ is the matrix with entries

$$M_{ij}(x) \stackrel{\text{def}}{=} \frac{\partial H_i}{\partial x_j}(x) \quad (34)$$

$$= \frac{1}{(d-1)!} \sum_{j_1, \dots, j_{d-1}=1}^n w_{i,j j_1 \dots j_{d-1}} x_{j_1} \dots x_{j_{d-1}} \quad (35)$$

that is

$$Z = \exp(-\text{tr} \log(I - M(G(y)))) \quad (36)$$

$$= \frac{1}{\det \left[\frac{\partial F}{\partial x}(G(y)) \right]} \quad (37)$$

$$= \frac{1}{JF(G(y))} . \quad (38)$$


In the case where $F_i(x) = x_i - H_i(x)$ with the $H_i(x)$ homogenous of the same degree d , it is easy to show that the Jacobian condition is equivalent to $M(x)$ being nilpotent for all x (see [10]). There are essentially two ways to express this

$$M(x)^n = 0, \quad \forall x \in \mathbb{C}^n \quad (39)$$

or

$$\text{tr} (M(x)^k) = 0, \quad \forall k \geq 1, \forall x \in \mathbb{C}^n \quad (40)$$


Equation (39) means that when one considers a *chain* (or caterpillar) diagram like



$$(41)$$

with n w -vertices, its contribution, for fixed i and j , is zero after *symmetrization* of the indices of the $n(d-1)$ incoming lower legs.

Equation (40) means that *loop* diagrams like



$$(42)$$

with $k \geq 1$ w -vertices, vanish after *symmetrization* of the indices of the $k(d-1)$ incoming legs. The formal inverse approach to the Jacobian conjecture can now be rephrased as the following

Problem : Show *explicitly* that in the polynomial algebra $\mathbb{C}[w]$ with indeterminates given by the tensor elements $w_{i,j_1\dots j_d}$, the $y = 0$ connected correlation functions

$$< \phi_i \bar{\phi}_{j_1} \dots \bar{\phi}_{j_N} >_{y=0}^c$$

belong to the *radical* of the ideal generated by the *symmetrized* chains and/or loops, provided the degree N is large enough.

This statement is by the Hilbert nullstellensatz *equivalent* to the Jacobian conjecture. It is even a theorem due to S. Wang [38] in the ($d = 2$) quadratic case. The proof is non constructive however, and an explicit combinatorial argument is an urgent desideratum.

III.2 Chains and/or loops?

Let $\mathfrak{c}$ be the ideal of $\mathbb{C}[w]$ generated by the symmetrized chains of length n , and let $\mathfrak{l}$ be the ideal generated by the symmetrized loops of length $k \geq 1$. While it is very tempting to work with $\mathfrak{c}$, it seems more fundamental to use $\mathfrak{l}$. This conclusion is implicit in [40]. Indeed the author uses the diagonal minor sums, i.e. the elementary symmetric functions of the eigenvalues, to express the nilpotence of $M(x)$, instead of the matrix elements of $M(x)^n$. We use the loops, that is the Newton power sums of the eigenvalues, which makes no difference since our ground ring is $\mathbb{C}$. Note that $\mathfrak{c} \subset \mathfrak{l}$: this is the Cayley-Hamilton theorem, i.e. “the Jacobian problem for $d = 1$ ”! But we also have $\mathfrak{l} \subset \sqrt{\mathfrak{c}}$, trivially because a nilpotent matrix must have zero eigenvalues and therefore the Newton sums of these eigenvalues are zero. It is very instructive to understand these two elementary statements in a purely combinatorial way. Regarding the first inclusion, we were surprised to find in the recent literature a combinatorial proof, with a flavor of loop-erased random walk, of the eminently classical Cayley-Hamilton theorem [37]. As for the second inclusion, there is a very nice explicit Fermionic proof [12] that, for a generic $n \times n$ matrix N , $(\text{tr } N)^{k(n-1)+1}$ is in the ideal generated by the matrix elements of N^k .

To see why the ideal $\mathfrak{c}$ is tempting to work with, we need to recall a theorem, first conjectured by Wang in the quadratic case [38], and proved in full generality by O. Gabber (see [10]).

Theorem 2 *If $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is globally invertible with polynomial inverse G then*

$$\deg G \leq (\deg F)^{n-1} \tag{43}$$

where $\deg F \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} (\deg F_i)$ and likewise for G .

In our context, this means that the vanishing of the connected correlation functions $\langle \phi_i \bar{\phi}_{j_1} \dots \bar{\phi}_{j_N} \rangle_{y=0}^c$ should happen as soon as $N > d^{n-1}$. Note that this bound is saturated by the well known triangular example given by $F_i(x) = x_i - x_{i+1}^d$, for $1 \leq i \leq n-1$, and $F_n(x) = x_n$. But d^{n-1} is the maximal number of leaves of those of our trees which have a depth less than or equal to $n-1$. If the chains in (41) needed not be symmetrized, the Jacobian conjecture would be trivial! Indeed, a tree with more than d^{n-1} leaves must have a chain of length at least n , going from the root ϕ_i to one of the leaves $\bar{\phi}_{j_\alpha}$. This observation, which goes back to [10], was likely the main impetus behind the formal inverse approach.

Remark that if we condition the sum over Feynman diagrams for the correlations $\langle \phi_i \bar{\phi}_{j_1} \dots \bar{\phi}_{j_N} \rangle_{y=0}^c$, by requiring the path between the root ϕ_i and a *specified* leaf $\bar{\phi}_{j_\alpha}$ to be of a certain length $\geq n$; the branches will be automatically symmetrized and the result would be zero. The problem is that we cannot know *in advance* which leaf will be linked to the root by a *long chain*.

In relation to previously used formal inversion formulas, let us mention that it is against QFT wisdom to mix the index space $\{1, \dots, n\}$ and the abstract space E that labels the vertices, as far as the combinatorics are concerned. From a QFT point of view, which admittedly is only one among many on the Jacobian conjecture, it is unnatural to use sums over colored or planar objects, as this reduces symmetry in the resulting expansion instead of enhancing it. We nevertheless concede the point that planarity can serve to “locate” the long chain, and order the trees accordingly, which is the main ingredient of the combinatorial “tour de force” of [34].

One of the cases treated in the latter article is that of $F_i(x) = x_i - H_i(x)$, with the H_i homogenous of the same degree d and the matrix $M(x)$ nilpotent of order 2. This has already been treated in [10] and [13] for instance, but let us sketch how to prove this result with our QFT model. The argument is adapted from an idea by V. Rivasseau.

First perform the translation change of variables $\phi \rightarrow \phi + y$, $\bar{\phi} \rightarrow \bar{\phi}$ in (3) to get, using $Z = 1$,

$$G_i(y) = y_i + \int d\mu(\bar{\phi}, \phi) \phi_i e^{\bar{\phi} H(\phi+y)} \quad (44)$$

where $d\mu(\bar{\phi}, \phi) \stackrel{\text{def}}{=} d\bar{\phi} d\phi e^{-\bar{\phi}\phi}$. This unorthodox change of variables used

in (44), which treats ϕ and $\bar{\phi}$ as independent variables and not as complex conjugates of one another, can be justified *a posteriori* by comparing the diagrammatic expansions on both sides of the equation. One can integrate the source ϕ_i by parts to get

$$G_i(y) = y_i + \int d\mu(\bar{\phi}, \phi) \frac{\partial}{\partial \bar{\phi}_i} e^{\bar{\phi}H(\phi+y)} \quad (45)$$

$$= y_i + \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) e^{\bar{\phi}H(\phi+y)} \quad (46)$$

$$= y_i + \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) e^{s\bar{\phi}H(\phi+y)} \Big|_{s=1} . \quad (47)$$

Then, interpolate between $s = 1$ and $s = 0$ to get

$$G_i(y) = y_i + \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) + \int_0^1 ds \Omega_i(s, y) \quad (48)$$

where

$$\Omega_i(s, y) \stackrel{\text{def}}{=} \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) [\bar{\phi}H(\phi+y)] e^{s\bar{\phi}H(\phi+y)} . \quad (49)$$

Notice that the second term of (48) reduces to $H_i(y)$, whereas for the third we have, by integrating the $\bar{\phi}$ by parts

$$\Omega_i(s, y) = \sum_{j=1}^n \int d\mu(\bar{\phi}, \phi) \frac{\partial}{\partial \phi_j} \left(H_i(\phi+y) H_j(\phi+y) e^{s\bar{\phi}H(\phi+y)} \right) \quad (50)$$

$$= \Omega_i^1(s, y) + \Omega_i^2(s, y) + \Omega_i^3(s, y) \quad (51)$$

where

$$\Omega_i^1(s, y) \stackrel{\text{def}}{=} \int d\mu(\bar{\phi}, \phi) \left(\sum_{j=1}^n M_{ij}(\phi+y) H_j(\phi+y) \right) e^{s\bar{\phi}H(\phi+y)} \quad (52)$$

$$\Omega_i^2(s, y) \stackrel{\text{def}}{=} \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) \left(\sum_{j=1}^n M_{jj}(\phi+y) \right) e^{s\bar{\phi}H(\phi+y)} \quad (53)$$

and

$$\Omega_i^3(s, y) \stackrel{\text{def}}{=} \int d\mu(\bar{\phi}, \phi) H_i(\phi+y) \left(\sum_{j,k=1}^n H_j(\phi+y) s \bar{\phi}_k M_{kj}(\phi+y) \right) e^{s\bar{\phi}H(\phi+y)} \quad (54)$$

Note that $\Omega_i^2(s, y) = 0$ since it contains $\sum_{j=1}^n M_{jj}(x) = \text{tr } M(x)$ at $x = \phi + y$, and $M(x)$ is nilpotent. Likewise, $\Omega_i^1(s, y)$ and $\Omega_i^3(s, y)$ vanish since the integrand contains a factor of the form

$$\sum_{j=1}^n M_{kj}(x) H_j(x) = \frac{1}{d} \sum_{j,l=1}^n M_{kj}(x) M_{jl}(x) x_l \quad (55)$$

$$= \frac{1}{d} \sum_{l=1}^n [M(x)^2]_{kl} x_l \quad (56)$$

$$= 0 \quad (57)$$

where we used Euler's identity for the homogenous H_i 's, and the fact that $M(x)$ is nilpotent of order 2. As a result $G_i(y) = y_i + H_i(y)$.

III.3 The Pauli exclusion principle

In order to be able to prove the Jacobian conjecture by purely combinatorial means, one needs to exhibit a volume effect similar to the Pauli exclusion principle, as otherwise one would not see the finiteness of the index set $\{1, \dots, n\}$ within the strictly tensorial Feynman diagrammatic notation were indices are contracted i.e. summed over. One would love to have Fermions, instead Bosons, entering the picture. Let us mention three, typically field theoretic, ideas that have not been pursued in previous attempts with the formal inverse approach, and which deserve further investigation.

III.3.1 Supersymmetry

One way to introduce Fermions in a purely Bosonic model is to exhibit a supersymmetry. If this could be done; it would probably be the “voie royale” towards understanding the conjecture. Unfortunately we have not been able to make much headway in this direction so far. Let us simply mention a strange feature of our model that hints towards a hidden supersymmetry. As a result of our choice of vertices and the fact that the propagators are directed, the perturbation expansion of the one-point function is reduced to a tree graph expansion. This means that the semi-classical expansion of our model around the “false vacuum” $\phi = 0, \bar{\phi} = 0$ is *exact*. Besides, the “integrals” in (3) which are supposed to be over $\mathbb{C}^n$ reduce to the contribution

of a single critical point: the “true vacuum” obtained by solving

$$\frac{\partial}{\partial \phi_i} (\bar{\phi} F(\phi) - \bar{\phi} y) = 0 \quad (58)$$


and

$$\frac{\partial}{\partial \bar{\phi}_i} (\bar{\phi} F(\phi) - \bar{\phi} y) = 0 \quad (59)$$

that is $\phi = G(y)$ and $\bar{\phi} = 0$. This is reminiscent of the Duistermaat-Heckman theorem [16] which is known to involve supersymmetry (see [39]).

III.3.2 Renormalization

The Gabber inverse degree bound, together with the previously given example that saturates it, suggest that the sought exclusion principle has to act along the chains from the root to the leaves of the trees but not across, i.e. within generations. This is quite odd in view of the eventual introduction of Fermionic variables in our model. This however hints to the possibility that the problem may come from “divergent” two-point subgraphs i.e. parts of the diagrams that look like



(60)

where the two indices i and j coincide. This leads to the following.


Question : Is there a way to eliminate these “divergent” pieces by adding, to the “action” $\bar{\phi} F(\phi) - \bar{\phi} y$, counter-terms that are made of symmetrized loops?

This is possible for $d = 1$, that is the Cayley-Hamilton theorem, but does not seem to be the consequence of a natural renormalization condition on the two-point correlation function. It would be interesting to explore this

idea using the new point of view on renormalization pioneered by A. Connes and D. Kreimer [14] since one of their motivations was the study of formal diffeomorphisms which is clearly related to our subject material.

III.3.3 Reverse Mayer expansion

We have repeatedly mentioned the Cayley-Hamilton theorem as the $d = 1$ case of the Jacobian conjecture. Let us now make this more precise. If $d = 1$ then $F_i(x) = x_i - H_i(x)$ with $H_i(x) = \sum_{j=1}^n w_{i,j} x_j$ and $M(x) = M = (w_{i,j})_{1 \leq i,j \leq n}$. When $y = 0$, the only interesting connected correlation functions $\langle \phi_i \phi_{j_1} \dots \phi_{j_N} \rangle_{y=0}^c$ are for $N = 1$, and their Feynman diagram expansion only generates chains of the form



(61)

If we use the loop ideal $\mathfrak{l}$ in the formulation of the problem in section III.1, then the ensuing statement that chains of length $\geq n$ belong to $\sqrt{\mathfrak{l}}$, or simply $\mathfrak{l}$ here, *is* the Cayley-Hamilton theorem. Therefore any combinatorial solution of this problem, for general d , should *at the very least* reproduce this well known result. Now the Cayley-Hamilton theorem is the statement that the derivative of $\frac{1}{Z} = \det(I - M)$, with respect to a matrix element of M , has a vanishing component in degree n in the M_{ij} variables. This in turn stems from the fact that $\frac{1}{Z}$ is itself a *polynomial* of degree n in M . We all know this from our “Fermionic/determinantal” upbringing, but let us suppose for a moment that all we know is Bosons and the only expression available to us is

$$Z = \int d\bar{\phi} d\phi e^{-\bar{\phi}\phi + \bar{\phi} M \phi} . \quad (62)$$

It is amusing to prove that $\frac{1}{Z}$ is a polynomial in M of degree at most n , using this formula. Let us explain an answer that is inspired from the Mayer expansion in statistical mechanics (see [33]), and constructive QFT (see [30]).

Consider the partition function $\mathcal{Z}$ of a gas of propagators


(63)

and vertices


$$\begin{array}{c} \leftarrow \\ i \end{array} \bullet \begin{array}{c} \leftarrow \\ j \end{array} = M_{ij} \quad (64)$$

with two kinds of interactions.

- The propagators attach to the vertices in all possible ways, as when applying Wick's theorem.
- The propagators can interact by *Mayer-links* represented by a squiggly line

$$\begin{array}{c} | \\ i \end{array} \text{---} \text{---} \text{---} \begin{array}{c} | \\ j \end{array} \quad (65)$$


and carrying a factor $-\delta_{ij}$ where i and j are the indices of the two propagators. One also assigns by hand a factor -1 per propagator-vertex loop, which we confess is cheating a bit. $\mathcal{Z}$ is therefore a sum of objects like



$$\quad (66)$$

with the appropriate symmetry factors, that is $\frac{1}{k}$ for each oriented loop with k vertices, an overall $\frac{1}{m!}$ if there are m loops, and a global sign $(-1)^m$. On the

one hand, if one sums over the structure of Mayer-links, with fixed contraction of the propagators to the vertices, one rebuilds a “hardcore constraint” factor $(1 - \delta_{ij})$ for each pair of propagators, which is the opposite of the operation one would do in a standard Mayer expansion. Since the available index space $\{1, \dots, n\}$ has cardinality n , there cannot be more than n propagators in a nonvanishing graph, ergo $\mathcal{Z}$ is of degree at most n in M . On the other hand, $\log \mathcal{Z}$, is a sum over connected objects like those of (66), where connectedness involves *both* types of lines. When one sums over Wick contractions, with fixed configuration of Mayer-links, the result is zero as soon as there is at least one Mayer-link, because of the following *exchange move* along the Mayer-link.


(67)

Indeed such a move does not affect the amplitudes, but modifies the loop count by one unit and thus the sign of the graph (compare with [37]). As a result, $\log \mathcal{Z}$ is a sum over single loops without Mayer-links and with a single (-1) factor. That is $\log \mathcal{Z} = -\log Z$ which concludes the argument.

An interesting question raised by this approach is

Question : Is there a hyperdeterminant, in the sense of [19], that would play, when $d \geq 2$, the role played by $\frac{1}{Z}$ when $d = 1$, and that would, upon derivation with respect to a tensor element $w_{i,j_1 \dots j_d}$ around a solution of the Jacobian condition, give some finiteness information on our correlation functions?

III.3.4 Are these three ideas different, really?

Although we have no precise unified framework to propose at the moment, we are tempted to say no. The interplay between supersymmetry, renormalization and the Mayer expansion is quite mysterious and is probably related to the combinatorics of the symmetric group and the inclusion-exclusion principle.

We will conclude by pointing out a few references where some clues on these relationships might be found. Mayer expansions involve coefficients which are Moebius functions of certain partition lattices (see [31]). These

coefficients can be calculated by an analog of the classical forest formula of Zimmermann in renormalization theory (see the introduction to chap. 4 of [3]). They can also be expressed using the so called Brydges-Kennedy forest formula [11] that was first proved there using the Hamilton-Jacobi equation. In [4] we gave a purely algebraic proof of the latter using some partial fraction combinatorial identities (Lemma II.2 in [4]). Such identities have been given a very elegant interpretation in terms of minimal factorizations of permutations as a product of transpositions [26]. The global sign in the Mayer coefficients is $(-1)^k$ where k is the number of edges in the forest. This sign obviously becomes the signature of the permutation in the latter interpretation. This strongly suggests a relationship between Mayer expansions and Fermions which was also alluded to in [30]. Note finally that the Brydges-Kennedy identity was considerably generalized in [5] (section III.2.1), where critical use is made of shuffles and a kind of Chen's lemma (see remarks following the proof of Lemma 9), although we did not know this at the time.

IV Conclusion

We hope to have provided enough evidence that the Jacobian conjecture is a very beautiful combinatorial challenge, where mathematicians, either conceptually or computationally inclined, and theoretical physicists could fruitfully share their knowledge. While future progress on the conjecture itself is still uncertain, there are bound to be benefits from such an interdisciplinary collaboration on this problem.

References

- [1] A. Abdesselam, *Feynman diagrams in algebraic combinatorics*, in preparation.
- [2] A. Abdesselam, *A physicist's proof of the Lagrange-Good multivariable inversion formula*, preprint, (2002).
- [3] A. Abdesselam, *Renormalisation constructive explicite*, Ph. D. Thesis, École Polytechnique, (1997).
- [4] A. Abdesselam and V. Rivasseau, *Trees, forests and jungles: a botanical garden for cluster expansions*. In: Constructive Physics, 7-36, V. Rivasseau, ed., Lecture Notes in Physics 446, Springer Verlag, 1995.

- [5] A. Abdesselam and V. Rivasseau, *An explicit large versus small field multiscale cluster expansion*, Rev. Math. Phys. Vol. **9** No 2, 123-199 (1997).
- [6] S. S. Abhyankar, *Historical ramblings in algebraic geometry and related algebra*, Amer. Math. Monthly **83** No 6, 409-448 (1976).
- [7] S. S. Abhyankar, *Lectures in algebraic geometry*, Notes by Chris Christensen, Purdue Univ. , 1974.
- [8] D. Bar-Natan, S. Garoufalidis, L. Rozansky and D. Thurston, *The Aarhus integral of rational homology 3-spheres I: A highly non trivial flat connection on S^3* , q-alg/9706004, preprint, (1997).
- [9] D. Bar-Natan, S. Garoufalidis, L. Rozansky and D. Thurston, *The Aarhus integral of rational homology 3-spheres II: Invariance and universality*, math.QA/9801049, preprint, (1998).
- [10] H. Bass, E. H. Connell and D. Wright, *The Jacobian conjecture: reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. (N.S.) **7** No 2, 287-330, (1982).
- [11] D. Brydges and T. Kennedy, *Mayer expansions and the Hamilton-Jacobi equation*, J. Stat. Phys. **48** No 1-2, 19-49, (1987).
- [12] C. de Calan and J. Magnen, personal communication, (2002).
- [13] C. Cheng, T. Sakkalis and S. Wang, *A case of the Jacobian conjecture*, J. Pure Appl. Algebra **96** No 1, 15-18, (1994).
- [14] A. Connes and D. Kreimer, *Hopf algebras, renormalization and noncommutative geometry*, Commun. Math. Phys. **199** No 1, 203-242, (1998).
- [15] G. Darboux, *Sur la série de Laplace*, C.R. Acad. Sci. Paris **68**, 324-327, (1869).
- [16] J. J. Duistermaat and G. J. Heckman, *On the variation in the cohomology of the symplectic form of the reduced phase space*, Invent. Math. **69** No 2, 259-268, (1982).
- [17] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics, 190, Basel, Birkhäuser verlag, 2000.
- [18] G. Gallavotti, *Perturbation Theory*, In: Mathematical physics towards the XXI century, 275-294, R. Sen and A. Gersten, eds., Ben Sheva, Ben Gurion University Press, 1994.

- [19] I. M. Gel'fand, M. M. Kapranov and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Mathematics: Theory & Applications, Boston MA, Birkhäuser Boston, Inc., 1994.
- [20] I. M. Gessel, *A combinatorial proof of the multivariable Lagrange inversion formula*, J. Combin. Theory Ser. A **45** No 2, 178-195, (1987).
- [21] I. J. Good, *Generalization to several variables of Lagrange's expansion, with applications to stochastic processes*, Proc. Cambridge Philos. Soc. **56**, 367-380, (1960).
- [22] P. Henrici, *Die Lagrange-Bürmannsche Formel bei formalen Potenzreihen*, Jahresber. Deutsch. Math. Verein. **86**, 115-134, (1984).
- [23] C. G. J. Jacobi, *De resolutione aequationum per series infinitas*, J. Reine Angew. Math. **6**, 257-286, (1830).
- [24] A. Joyal, *Une théorie combinatoire des séries formelles*, Adv. Math **42**, 1-82, (1981).
- [25] O. H. Keller, *Ganze Cremona-Transformationen*, Monats. Math. Physik **47**, 299-306 (1939).
- [26] V. Lafforgue, letter to the author, (1994).
- [27] J. L. de Lagrange, *Nouvelle méthode pour résoudre des équations littérales par le moyen des séries*, Mém. Acad. Roy. Sci. Belles-Lettres Berlin **24** (1770).
- [28] P. S. Laplace, *Mémoire sur l'usage du calcul aux différences partielles dans la théorie des suites*, Mém. Acad. Roy. Sci. Paris 99-122, (1777).
- [29] H. Poincaré, *Sur les résidus des intégrales doubles*, Acta. Math. **9**, 321-380, (1886).
- [30] V. Rivasseau, *From perturbative to constructive renormalization*, Princeton NJ, Princeton University Press, 1991.
- [31] G.-C. Rota, *On the foundations of combinatorial theory I. Theory of Möbius functions*, Z. Wahrscheinlichkeitstheorie **2**, 340-368, (1964).
- [32] L. Rozansky, *A universal $U(1)$ -RCC invariant of links and rationality conjecture*, math.GT/0201139, preprint, (2002).
- [33] D. Ruelle, *Statistical mechanics: Rigorous results*, New York-Amsterdam, W. A. Benjamin, 1969.
- [34] D. Singer, *On Catalan trees and the Jacobian conjecture*, Electron. J. Combin. **8** No 1, Research paper 2, 35 pp., (2001).

- [35] S. Smale, *Mathematical problems for the next century*, Math. Intelligencer **20** No 2, 7 (1998).
- [36] T. J. Stieltjes, *Sur une généralisation de la série de Lagrange*, Ann. Sci. École Normale. Paris Sér. 3 **2**, 93-98, (1885).
- [37] H. Straubing, *A combinatorial proof of the Cayley-Hamilton theorem*, Discrete Math. **43** No 2-3, 273-279, (1983).
- [38] S. Wang, *A Jacobian criterion for separability*, J. Algebra **65**, 453-494, (1980).
- [39] E. Witten, *Two-dimensional gauge theories revisited*, J. Geom. Phys. **9** No 4, 303-368, (1992).
- [40] D. Wright, *Reversion, trees, and the Jacobian conjecture*. In: Combinatorial and computational algebra, 249-267, Contemp. Math., 264, Providence RI, Amer. Math. Soc., 2000.
- [41] D. Wright, *The tree formulas for reversion of power series*, J. Pure Appl. Algebra **57** No 2, 191-211, (1989).
- [42] J. Zinn-Justin, *Quantum field theory and critical phenomena*, New York, The Clarendon Press, Oxford University Press, 1989.